

***AGRICULTURAL LAND CLASSIFICATION***

**Pegasus Group**

*Nuneham Estate*



**Our Ref: SES/PG/NE/#1**

**Date: 11<sup>th</sup> Oct 2022**

**Client:**

Pegasus Group  
3<sup>rd</sup> Floor, Gainsborough House  
30-40 Grey Street  
Newcastle upon Tyne  
NE1 6AE

***AGRICULTURAL LAND CLASSIFICATION***

***Nuneham Estate***

A report prepared on behalf of *Soil Environment Services* by:

**Dr Robin S Davies** BSc PhD F.I.SoilSci PGC *Contaminated Land Management*  
*Managing Director*

*This report has been prepared by Soil Environment Services with all reasonable skill, care and diligence, within the terms of The Contract with The Client. The report is the property of The Client who can assign this report to any third party who will then be afforded the same assurances as detailed within the terms of the original Contract with The Client.*

---

**Soil Environment Services**

Agricultural Land Classification, Contaminated Land  
Risk Assessment, Mineral Extraction Soil Planning  
Unit 8, Stocksfield Hill, Stocksfield, Northumberland, NE43 7TN  
Tel: 01661 844 827, Email: rd@soilenvironmentservices.co.uk  
[www.soilenvironmentservices.co.uk](http://www.soilenvironmentservices.co.uk)

---

# CONTENTS

	<b>Page</b>
<b>1. INTRODUCTION</b>	
<b>1.1 Methodology</b>	<b>4</b>
<b>1.2 Previous ALC gradings</b>	<b>4</b>
<b>2. CLIMATIC LIMITATIONS</b>	<b>5</b>
<b>2.1 Overall climate</b>	<b>5</b>
<b>2.2 Local climate</b>	<b>5</b>
<b>3. SITE LIMITATIONS</b>	<b>5</b>
<b>3.1 Gradient</b>	<b>5</b>
<b>3.2 Microrelief</b>	<b>5</b>
<b>3.3 Flooding</b>	<b>6</b>
<b>4. SOIL LIMITATIONS</b>	<b>7</b>
<b>4.1 Texture and structure</b>	<b>7</b>
<b>4.2 Depth</b>	<b>7</b>
<b>4.3 Stoniness</b>	<b>7</b>
<b>4.4 Chemical</b>	<b>7</b>
<b>5. INTERACTIVE LIMITATIONS</b>	<b>8</b>
<b>5.1 Wetness</b>	<b>8</b>
<b>5.2 Droughtiness</b>	<b>8</b>
<b>5.3 Erosion</b>	<b>8</b>
<b>6. AGRICULTURAL LAND CLASSIFICATION</b>	<b>9</b>
<b>6.1 Most limiting factor</b>	<b>9</b>
<b>6.2 Current ALC grading</b>	<b>9</b>

**DRAWING 1**            **ALC Grade and survey points**

**APPENDIX A**            **Survey profile data sheet**

**STATEMENT OF COMPETENCE**

**GENERAL INFORMATION SOURCES**

**GLOSSARY**

## 1. INTRODUCTION

An Agricultural Land Classification (ALC) has been carried out on ~57 ha of land on the Nuneham Estate (Drawing 1). The site is centred on OS Grid Ref. 454358,200060.

The survey was conducted on the 22<sup>nd</sup> October 2022 and classified the land into one or more of the below grades (see Drawing 1). On the survey date, the site was in agricultural use.

### 1.1 Methodology

Agricultural land is classified into the following grades according to the 1988 guidelines<sup>1</sup>.

Grade	Description
1	<b>Excellent quality agricultural land</b> with no or very minor limitations to agricultural use.
2	<b>Very good quality agricultural land</b> with minor limitations which affect crop yield, cultivation or harvesting.
3a	<b>Good quality agricultural land</b> capable of producing moderate to high yields of a narrow range of arable crops or moderate yields of a wider range of crops.
3b	<b>Moderate quality agricultural land</b> capable of producing moderate yields of a narrow range of crops or lower yields of a wider range of crops.
4	<b>Poor quality agricultural land</b> with severe limitations which significantly restrict the range of crops and/or level of yields.
5	<b>Very poor quality agricultural land</b> with very severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

The classification includes an initial desktop investigation to examine previously mapped soil types and to note the drift and solid geology followed by the field survey consisting of auger borings at one every 100 m in general and a pit excavated in each of the main soil types to confirm the structures and stone content if needed. Laboratory analysis of soil textures is undertaken if needed in order to confirm textures such the *heavy/medium* clay and *medium/fine* sand categories or stone content. All site survey profile data is listed in Appendix A.

All of the potential limitations are assessed and then the most limiting factor dictating the ALC grade was determined for this site and is detailed in Table 2.

### 1.2 Previous ALC gradings

Grading on the MAFF (1983) 1:250 000 Provisional map indicated the site is located on ALC Grade 3 land with Grade 4 just on the west. A detailed survey has not previously been undertaken.

## 2. CLIMATIC LIMITATIONS

### 2.1 Overall climate

The climatological data for the site centre is detailed in Table 1.

<b>Table 1</b>		
<b>Climatological information<sup>3</sup></b>		
<b>Factor</b>	<b>Units</b>	<b>Value</b>
Altitude AOD	m	60
Accumulated temperature	day°C (Jan-June)	1471.1
Average Annual Rainfall	mm	838.8
Field Capacity Days	days	185.8
Moisture Deficit Wheat	mm	95.2
Moisture Deficit Potatoes	mm	84.2
Overall climate ALC Grade	Grade 2	

Overall climate will not result in the most significant limiting factor for this site.

### 2.2. Local climate

Local climate will not result in a significant limiting factor for this site.

## 3 SITE LIMITATIONS

### 3.1 Gradient

The gradient will not result in a significant limiting factor for this site.

### 3.2 Microrelief

The microrelief will not result in a significant limiting factor for this site.

### 3.3 Flooding

A low risk of flooding from surface water in places has been identified across the site (<https://flood-warning-information.service.gov.uk/long-term-flood-risk>).



## 4 SOIL LIMITATIONS

### 4.1 Texture and structure

The soils in general are mostly heavy clay loam topsoils over clay subsoils with the subsoil structure of coarse angular blocky above 60 cm depth. Only slight variations exist across most of the site. Calcareous topsoils were only found in the north east. In the far south east, the soils are sandy loams over sand.

The site has previously been mapped as having mainly soils of the following Association/s.

Most of the site - Wickham 2: *Slowly permeable seasonally waterlogged fine loamy over clayey, fine silty over clayey and clayey soils. Small areas of slowly permeable calcareous soils on steeper slopes.*

Far south east corner – Frilford: *Deep well drained sandy and coarse loamy soils. Some ferruginous sandy and some coarse loamy soils affected by groundwater. Risk of water erosion.*

([www.landis.org.uk](http://www.landis.org.uk)).

### Superficial Geology – 1:50 000 scales

*None recorded over most of the site and Head - Clay, silt, sand and gravel. Sedimentary superficial deposit in the north western area.*

### Bedrock Geology - 1:50 000 scale bedrock geology description

Most of the site: *Amphill Clay Formation and Kimmeridge Clay Formation - Mudstone. Sedimentary bedrock*

South east corner: *Kimmeridge Clay Formation - Siltstone and sandstone and Portland Group - Limestone and calcareous sandstone.*

### 4.2 Depth

Soil depth will not result in a significant limiting factor for this site.

### 4.3 Stoniness

Stoniness is not a direct significant limiting factor for soils noted on site but does contribute to the droughtiness.

### 4.4 Chemical

Chemical contamination is not a significant limiting factor for this site

## **5. INTERACTIVE LIMITATIONS**

### **5.1 Wetness**

Mottling was noted at shallow depths below the topsoil across most of the site and a slowly permeable subsoil existed in general above 60 cm depth. The combination of a Wetness Class of III for the soils (see Appendix A) with the Field Capacity Days of 127.8 and a topsoil texture of heavy clay loam results in a wetness limitation and hence ALC Grade 3b or 3a where calcareous in the north east.

### **5.2. Droughtiness**

The Available Water Capacity which subsequently when considered with respect to the Moisture Deficit for wheat and potatoes results in a Moisture Balance giving a main limitation for both wheat and potatoes on the sandy soils in the south east. This results in an ALC Grade of 2.

### **5.3 Erosion**

Erosion will not result in a significant limiting factor for this site.



## 6. AGRICULTURAL LAND CLASSIFICATION

### 6.1 Most limiting factor/s

#### *Grade 3b*

A Wetness Limitation for the majority of the site with the clayey non-calcareous topsoils.

#### *Grade 3b*

A Wetness Limitation for the clayey soils with calcareous topsoils in the north east..

#### *Grade 2*

A Droughtiness Limitation for the sandy soils in the south east.

### 6.2 Current grading

This survey has resulted in an Agricultural Land Classification of the following grades (Drawing 1):

<b>Table 2. ALC gradings and limitations</b>			
<b>Grade</b>	<b>ha</b>	<b>%</b>	<b>Limitation</b>
<b>1</b>			
<b>2</b>	4	7.0	Droughtiness
<b>3a</b>	4	7.0	Wetness
<b>3b</b>	49	86.0	Wetness
<b>4</b>			
<b>5</b>			
<b>Non-agricultural land</b>			
<b>Total</b>	<b>57</b>	<b>100%</b>	

# **DRAWING 1**

**ALC Grade**

**Key**

ALC Grades

- Grade 1
- Grade 2
- Grade 3a
- Grade 3b
- Grade 4
- Grade 5
- Non agricultural land

- Boring
- Pit


# Soil Environment Services

Drawing Title: ALC Grade

Drawing No.: 1

Scale: 1:10000

Date: 07/07/2022



Ordnance Survey copy licence PMR0046161

# APPENDIX A

## Soil profile data

### Notes

1. All abbreviations relating to soil parameters are standard and derived from the guidance documents:  
  
*Agricultural Land Classification of England and Wales*. Revised guidelines and criteria for grading the quality of agricultural land. MAFF. 1988.  
*Soil Survey Field Handbook*. Technical Monograph No.5. Soil Survey of England and Wales.1976.
2. The pit data is detailed in this table and information on structure and stone content copied to the appropriate boring profiles.
3. Any blanks or zeros in the cells indicate the data is not needed or appropriate for that cell.
4. If 'NA' is inserted in a cell the information is not appropriate on this occasion.
5. Boring or pit locations are directly (within 2 m accuracy) on the grid reference corresponding to the points on the map unless otherwise stated.
6. A point directly marked on a track, boundary or other feature will be moved 2-3 m off the point or omitted if surrounding points and soil types allow.
7. Borings that are potentially within 15 m of a gas pipeline are limited to 0.4 m depth and the strata description in the data table below this depth will be extrapolated from nearby borings and upper strata characteristics.
8. The *Observation Density* is 1 per ha on a 100 m grid using a semi *Free Survey* method if appropriate\*. The letter 'B' in the second column of the data table refers to an observation point at which a boring may have been undertaken. In some situations it is not possible to visit the location due to for example crop status or animals in a field. In some cases the location is visited and observation of the soils at the surface is sufficient. In all cases the soil, geology, topography, flood risk and aerial crop patterns are assessed from published sources and the soils will be subject to a full 120 cm depth boring either side of a non-visited or non-bored point. If all data sources are agreeable, a soil pattern can be established.  
  
\* British Society of Soil Science. Working With Soil – The Professional Competency Scheme. Agricultural Land Classification: England and Wales. How2 sheet 4.2.4. 2018.
9. For moisture balance calculations, *strongly*, *moderately* and *well developed* structure will equate to *good*, *moderate* or *poor* structure terms respectively in Table 14 of the guidelines.
10. Pit information in addition to that listed in the table below will be detailed in Section 4.1 and 4.3 if needed.



Obs point	Grid ref. if off intersection	Boring or Pit	Grad. (deg)	Base Depth (cm)	OFFICE USE	Text.	Calc	Matrix colour	Mots./ black ferro. conc. %/ depth	Mott colour or FC if ferro. conc.	Ped face colour	Stns %	Stns type	Porosity	Struct (Firm consistency)	Degree of development	SPL depth (cm)	Gleying depth (cm)	SWC	Grade (wetness)	TAV	EAV	StTAV	StEAV	MBW	Grade (Drought: W/EART)	MBP	Grade (Drought: POTATOES)				
21		B	s7	30		HCL	N	10YR43				2	HR											18		1						
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
				120		C		10YR53	15/60	10YR56		0		P	CAB	MD								16	8	1	0.5					
22		B	s7	120								0												0	0	0	0					
				30		HCL	N	10YR43				2	HR													18		1				
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
23		B	s7	120		C		10YR53	15/60	10YR56		0		P	CAB	MD									0	0	0	0				
				30		HCL	N	10YR42				2	HR													18		1				
				60		C		10YR53	20/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	26.60	2	12.26	1	
24		B	s7	90		C		2.5Y52	10/60	10YR56		0	HR	P	CAB	MD									16	8	1	0.5				
				120		C		2.5Y51	20/90	7.5YR56		0			M											13	7	0	0			
				30		HCL	N	10YR42				2	HR														18		1			
25		B	s7	60		C		10YR53	20/30	10YR56		2	HR	P	CSAB	MD									16	8	1	0.5	26.60	2		
				90		C		2.5Y52	10/60	10YR56		0	HR	P	CAB	MD										16	8	1	0.5			
				120		C		2.5Y51	20/90	7.5YR56		0			M											13	7	0	0			
26		B	s7	30		HCL	N	10YR42				2	HR												18		1					
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	26.60	2	12.26	1	
				120		C		10YR53	15/60	10YR56		0		P	CAB	MD											16	8	1	0.5		
27		P	s7	120								0													0	0	0	0				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
28		B	s7	120		C		10YR53	15/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
29		B	s7	120		C		10YR53	15/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR42				2	HR														18		1			
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
30		B	s7	120		C		10YR53	15/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	20/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	26.60	2	12.26	1	
31		B	s7	90		C		2.5Y52	10/60	10YR56		0	HR	P	CAB	MD									16	8	1	0.5				
				120		C		2.5Y51	20/90	7.5YR56		0			M											13	7	0	0			
				30		HCL	N	10YR42				2	HR														18		1			
32		B	s7	60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD									16	8	1	0.5	26.60	2		
				120		C		10YR53	15/60	10YR56		0		P	CAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
				120								0															0	0	0	0		
33		B	s7	30		HCL	N	10YR43				2	HR												18		1					
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
				120		C		10YR53	15/60	10YR56		0		P	CAB	MD											16	8	1	0.5		
34		B	s7	120								0													0	0	0	0				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
35		B	s7	120		C		10YR53	10/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	10/30	10YR56		2		P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
36		B	s7	120		C		10YR53	10/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	10/30	10YR56		2		P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
37		B	s7	120		C		10YR53	10/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	10/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
38		B	s7	120		C		10YR53	15/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	15/30	10YR56		2	HR	P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
39		B	s7	120		C		10YR53	10/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	10/30	10YR56		2		P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
40		B	s7	120		C		10YR53	10/60	10YR56		0		P	CAB	MD									16	8	1	0.5				
				30		HCL	N	10YR43				2	HR														18		1			
				60		C		10YR53	10/30	10YR56		2		P	CSAB	MD			60	30	III	3b		16	8	1	0.5	29.60	2	12.26	1	
120		C		10YR53	10/60	10YR56		0		P	CAB	MD												16	8	1	0.5					
120												0												0	0	0	0					



SES Ltd undertake several dozen Agricultural Land Classification (ALC) or Land Capability Classifications for Agriculture (LCCA-Scotland) surveys a year and have worked on sites up to 1000 ha including housing, roads, solar farm and mineral extraction developments.. We have been undertaking ALC surveys for 25 years and have won many contracts to supply Land Classification reports to local authorities as part of their strategic development plans. A number of our staff have attended the training course Agricultural Land Classification: England and Wales. Working with Soil – The IPSS Professional Competency Scheme. BSSS & DEFRA.

### **DR ROBIN DAVIES BSc PhD F.I.SoilSci. (Managing Director)**

- Fellow of The British Society of Soil Science
- Council Member of The Institute of Professional Soil Scientists for 4 years.
- PhD Soil Physics - Agricultural land drainage - University of Newcastle upon Tyne
- Founder and Managing Director of Soil Environment Services Limited for 25 years.

#### *Selected peer reviewed scientific papers:*

- \* **Soil nitrogen depletion - the threat from soil stockpiling.** Environmental Scientist: Journal of The Institution of Environmental Sciences, 1997.
- \* **Nitrogen loss from a soil, restored after surface-mining.** Journal of Environmental Quality, 1995
- \* **The influence of soil factors on the growth of a grass/clover sward on a restored site in Northumberland.** Grass & Forage Science, 1994.
- \* **The effect of post-restoration cropping regime on some physical properties of a restored soil.** Soil Use & Management, 1994
- \* **Water availability in a restored soil.** Soil Use & Management, 1992.
- \* **A laboratory Method for Investigating the Stabilisation of Mole Channels.** J.Agric.Eng.Res.1991.

### **LOUISE TAVASSO BSc (Hons) . (Soil surveyor/ Environmental Consultant)**

**Member of** British Society of Soil Science  
**Postgraduate short course** Contaminated Land Risk assessment – LQM Nottingham University

Worked for Soil Environment Services Limited for 16 years. Environmental consultant with initial work in contaminated land risk assessment and since 2011 as assistant soil surveyor with last three years as lead consultant on agricultural land classification surveys. All work areas have required field survey and identification and description of soils combined with an understanding of soil processes for reporting.

Completed the BSSS Agricultural Land Classification Course – 2021.



#### *Main areas of specialisation*

##### **1 Agricultural Land Classification**

Soil survey and Agricultural Land Classification for planning applications –, roads, housing, solar parks. Fully conversant with the procedures of the *Agricultural Land Classification of England and Wales, Guidelines and criteria for grading the quality of agricultural land*, 1988, MAFF, London.

##### **2 Soil survey for habitat restoration**

Soil survey and nutrient analysis assessment for conversion of farmland to species rich grassland.

##### **3 Contaminated land risk assessment**

Phase 1 site survey risk assessment of contaminated land; site investigation, on-site monitoring; risk analysis, modelling and communication; recommendations for Phase 2 and remediation options.

#### *Examples of Agricultural Land Classification (ALC or LCCA Scotland) consultancy work*

Kier Mining. Greenburn Opencast Coal Site. Soils and deep peat survey for LCCA report soil resources planning. 2011

Newcastle International Airport Ltd. ALC survey for solar park development. 2021.

#### *Examples of soil survey habitat creation consultancy work*

BSG Ecology. Backwork Estate – farmland conversion to wildflower meadow. 2020.

Private garden owner. Soil survey and recommendation for drainage system design. 2021

#### *Examples of contaminated land consultancy work*

Numerous risk assessments on petrol stations for hydrocarbon leakages (2006-2019)

Farm building risk assessments for conversion to residential housing (2006-2019)



## GENERAL INFORMATION SOURCES

1. *Agricultural Land Classification of England and Wales*. Revised guidelines and criteria for grading the quality of agricultural land. MAFF. 1988.
2. *Soil Survey Field Handbook*. Technical Monograph No.5. Soil Survey of England and Wales.1976.
3. *Climatological Data for Agricultural Land Classification*, The Met. Office 1989
4. *Soil Map of England and Wales: 1:250 000*. Soil Survey of England and Wales, Harpenden.
5. *Soils and Their Use in South East England*. Soil Survey of England and Wales,
6. *Agricultural Land Classification Map* 1:250 000. MAFF 1983.
7. *Risk of Flooding*: <https://flood-warning-information.service.gov.uk/long-term-flood-risk>
8. *Geology of Britain Viewer*. Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved
9. *Butler, B E. Soil Classification for Soil Survey Monographs on Soil Survey (1980)* Clarendon Press, Oxford
10. *Munsell Soil Colour Charts, Munsell Colour, Grand Rapids 1994.*

# **GLOSSARY**

## ABBREVIATIONS AND TERMS USED IN SURVEY DATA

Soil pit and auger boring information collected during ALC survey is held on a computer database and is reproduced in this report. Terms used and abbreviations are set out below. These conform to definitions contained in the Soil Survey Field Handbook (Hodgson, 1997).

### 1. Terms used on computer database, in order of occurrence.

**GRID REF:** National 100 km grid square and 8 figure grid reference.

**LAND USE:** At the time of survey

<b>WHT:</b>	Wheat	<b>SBT:</b>	Sugar Beet	<b>HTH:</b>	Heathland
<b>BAR:</b>	Barley	<b>BRA:</b>	Brassicas	<b>BOG:</b>	Bog or Marsh
<b>OAT:</b>	Oats	<b>FCD:</b>	Fodder Crops	<b>DCW:</b>	Deciduous Wood
<b>CER:</b>	Cereals	<b>FRT:</b>	Soft and Top Fruit	<b>CFW:</b>	Coniferous Woodland
<b>MZE:</b>	Maize	<b>HRT:</b>	Horticultural Crops	<b>PLO:</b>	Ploughed
<b>OSR:</b>	Oilseed Rape	<b>LEY:</b>	Ley Grass	<b>FLW:</b>	Fallow (inc. Set aside)
<b>POT:</b>	Potatoes	<b>PGR:</b>	Permanent Pasture	<b>SAS:</b>	Set Aside (where known)
<b>LIN:</b>	Linseed	<b>RGR:</b>	Rough Grazing	<b>OTH:</b>	Other
<b>BEN:</b>	Field Beans	<b>SCR:</b>	Scrub		

**GRDNT:** Gradient as estimated or measured by hand-held optical clinometer.

**GLEYS, SPL:** Depth in centimetres to gleying or slowly permeable layer.

**AP (WHEAT/POTS):** Crop-adjusted available water capacity.

**MB (WHEAT/POTS):** Moisture Balance. (Crop adjusted AP - crop potential MD)

**DRT:** Best grade according to soil droughtiness.

If any of the following factors are considered significant, 'Y' will be entered in the relevant column.

<b>MREL:</b>	Microrelief limitation	<b>FLOOD:</b>	Flood risk	<b>EROSN:</b>	Soil erosion risk
<b>EXP:</b>	Exposure limitation	<b>FROST:</b>	Frost prone	<b>DIST:</b>	Disturbed land
<b>CHEM:</b>	Chemical limitation				

**LIMIT:** The main limitation to land quality: The following abbreviations are used.

<b>OC:</b>	Overall Climate	<b>AE:</b>	Aspect	<b>EX:</b>	Exposure
<b>FR:</b>	Frost Risk	<b>GR:</b>	Gradient	<b>MR:</b>	Microrelief
<b>FL:</b>	Flood Risk	<b>TX:</b>	Topsoil Texture	<b>DP:</b>	Soil Depth
<b>CH:</b>	Chemical	<b>WE:</b>	Wetness	<b>WK:</b>	Workability
<b>DR:</b>	Drought	<b>ER:</b>	Erosion Risk	<b>WD:</b>	Soil Wetness/Droughtiness

**ST:** Topsoil Stoniness

**TEXTURE:** Soil texture classes are denoted by the following abbreviations:-

<b>S:</b> Sand	<b>LS:</b> Loamy Sand	<b>SL:</b> Sandy Loam
<b>SZL:</b> Sandy Silt Loam	<b>CL:</b> Clay Loam	<b>ZCL:</b> Silty Clay Loam
<b>ZL:</b> Silt Loam	<b>SCL:</b> Sandy Clay Loam	<b>C:</b> Clay
<b>SC:</b> Sandy clay	<b>ZC:</b> Silty clay	<b>OL:</b> Organic Loam
<b>P:</b> Peat	<b>SP:</b> Sandy Peat	<b>LP:</b> Loamy Peat
<b>PL:</b> Peaty Loam	<b>PS:</b> Peaty Sand	<b>MZ:</b> Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of the following prefixes:-

**F:** Fine (more than 66% of the sand less than 0.2mm)  
**M:** Medium (less than 66% fine sand and less than 33% coarse sand)  
**C:** Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content: **M:** Medium (< 27% clay) **H:** heavy (27 - 35% clay)

**MOTTLE COL:** Mottle colour using Munsell notation.

**MOTTLE ABUN:** Mottle abundance, expressed as a percentage of the matrix or surface described.

**F:** few <2% **C:** common 2 - 20% **M:** many 20 - 40% **VM:** very many 40%+

**MOTTLE CONT:** Mottle contrast

**F:** faint - indistinct mottles, evident only on close inspection  
**D:** distinct - mottles are readily seen  
**P:** Prominent - mottling is conspicuous and one of the outstanding features of the horizon.

**PED. COL:** Ped face colour using Munsell notation.

**GLEYS:** If the soil horizon is gleyed a 'Y' will appear in this column. If slightly gleyed, an 'S' will appear.

**STONE LITH:** Stone Lithology - One of the following is used.

<b>HR:</b> All hard rocks and stones	<b>SLST:</b> Soft oolitic or dolimitic limestone
<b>CH:</b> Chalk	<b>FSST:</b> Soft, fine grained sandstone
<b>ZR:</b> Soft, argillaceous, or silty rocks	<b>GH:</b> Gravel with non-porous (hard) stones
<b>MSST:</b> Soft, medium grained sandstone	<b>GS:</b> Gravel with porous (soft) stones
<b>SI:</b> Soft weathered igneous or metamorphic rock	

Stone contents are given in % by volume for sizes >2cm, >6cm and total stone >2mm.

**STRUCT:** The degree of development, size and shape of soil peds are described using the following notation

<b><u>Degree of development</u></b>	<b>WA:</b> Weakly developed Adherent	<b>WK:</b> Weakly developed
	<b>MD:</b> Moderately developed	<b>ST:</b> Strongly developed
<b><u>Ped size</u></b>	<b>F:</b> Fine	<b>M:</b> Medium
	<b>C:</b> Coarse	<b>VC:</b> Very coarse
<b><u>Ped Shape</u></b>	<b>S:</b> Single grain	<b>M:</b> Massive
	<b>GR:</b> Granular	<b>AB:</b> Angular blocky
	<b>SAB:</b> Sub-angular blocky	<b>PR:</b> Prismatic
	<b>PL:</b> Platy	

**CONSIST:** Soil consistence is described using the following notation:

**L:** Loose      **VF:** Very Friable      **FR:** Friable      **FM:** Firm  
**VM:** Very firm      **EM:** Extremely firm      **EH:** Extremely Hard

**SUBS STR:** Subsoil structural condition recorded for the purpose of calculating profile droughtiness: **G:** Good      **M:** Moderate      **P:** Poor

**POR:** Soil porosity. If a soil horizon has poor porosity with less than 0.5% biopores >0.5mm, a 'Y' will appear in this column.

**IMP:** If the profile is impenetrable to rooting a 'Y' will appear in this column at the appropriate horizon.

**SPL:** Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.

**CALC:** If the soil horizon is calcareous with naturally occurring calcium carbonate exceeding 1% a 'Y' will appear this column.

## 2. Additional terms and abbreviations used mainly in soil pit descriptions.

**STONE ASSESSMENT:**

**V:** Visual      **S:** Sieved      **D:** Displacement

**MOTTLE SIZE:**

<b>EF:</b> Extremely fine <1mm	<b>M:</b> Medium 5-15mm
<b>VF:</b> Very fine 1-2mm>	<b>C:</b> Coarse >15mm
<b>F:</b> Fine 2-5mm	

**MOTTLE COLOUR:** May be described by Munsell notation or as ochreous (OM) or grey (GM).

**ROOT CHANNELS:** In topsoil the presence of 'rusty root channels' might also be noted as RRC.

**MANGANESE CONCRETIONS:** Assessed by volume

<b>N:</b> None	<b>M:</b> Many	20-40%
<b>F:</b> Few <2%	<b>VM:</b> Very Many	>40%
<b>C:</b> Common 2-20%		

**POROSITY:**

**P:** Poor - less than 0.5% biopores at least 0.5mm in diameter  
**G:** Good - more than 0.5% biopores at least 0.5mm in diameter

**ROOT ABUNDANCE:**

The number of roots per 100cm <sup>2</sup> :		Very Fine and Fine	Medium and Coarse
<b>F:</b>	Few	1-10	1 or 2
<b>C:</b>	Common	10.25	2 - 5
<b>M:</b>	Many	25-200	>5
<b>A:</b>	Abundant	>200	

**ROOT SIZE**

<b>VF:</b> Very fine <1mm	<b>M:</b> Medium	2 - 5mm
<b>F:</b> Fine 1-2mm	<b>C:</b> Coarse	>5mm

**HORIZON BOUNDARY DISTINCTNESS:**

<b>Sharp:</b> <0.5cm	<b>Gradual:</b> 6 - 13cm
<b>Abrupt:</b> 0.5 - 2.5cm	<b>Diffuse:</b> >13cm
<b>Clear:</b> 2.5 - 6cm	

**HORIZON BOUNDARY FORM:** Smooth, wavy, irregular or broken.\*

\* See Soil Survey Field Handbook (Hodgson, 1997) for details.